Sep. 1, 1981

[54] SPIROPYRAN COMPOUNDS OF PIPERIDINE OR THIAZINE RINGS

[75] Inventors: Robert J. Guglielmetti, Brest; Francis
Garnier, Champigny; Yves M.
Poirier, Brest; Gisele M. C. Petillon,

Briec, all of France

[73] Assignee: Etat Français represente par le

Delegue General pour l'Armement,

Paris, France

[21] Appl. No.: 96,395

[22] Filed: Nov. 21, 1979

[30] Foreign Application Priority Data

[51] Int. Cl.³ C07D 515/10; C07D 491/107 [52] U.S. Cl. 544/6; 546/17;

[56] References Cited

U.S. PATENT DOCUMENTS


OTHER PUBLICATIONS

Maguet et al., J. Heterocyclic Chem. vol. 15, Dec. 1978, pp. 1439-1446.

Primary Examiner—Henry R. Jiles
Assistant Examiner—Bernard Dentz
Attorney, Agent, or Firm—Stevens, Davis, Miller &
Mosher

[57] ABSTRACT

The invention relates to spiropyran compounds. They have the general formula (I):



in which:

X represents the CH₂ group or a sulfur atom; R₁ and R₂ represent electron donor and/or acceptor groups selected for instance from the group NO₂, COC₆H₅, CN, the halogens, the acyl substituents, alkoxy substituents, the hydrogen atom, the alkyl and mercaptoalkyl groups, the ethers of the formula CH₂OR, thioethers of the formula CH₂SR and amines of the formula CH₂NR₂ in which R represents an alkyl group; R₃ is a linear or branched alkyl group, having 1 to 18 carbon atoms, or an arylalkyl or alkenyl group.

 $-R_{4}$, R_{5} , R_{6} , R_{7} , R_{8} and R_{9} have one of the meanings indicated for R_{3} or represent, two by two, with the group in ortho position, a $-(CH_{2})_{n}$ — group in which

n is a number between 5 and 10.

 R_{10} represents a hydrogen atom if X=S and if $X=CH_2$ a hydrogen atom, an alkyl group having 1 to 18 carbon atoms, an arylalkyl group in which the alkyl group has 1 to 5 carbon atoms, or an alkenyl group having 2 to 10 carbon atoms.

The compounds in accordance with the invention have photochromic properties and are suitable for the recording of information on transparent or opaque supports.

6 Claims, No Drawings